



*COLLEGE ON SOIL PHYSICS*

# ARIDITY AND DROUGHT INDICES

**Donald Gabriels**

**Centre for Eremology, Ghent University, Belgium  
Scientific Consultant CAZALAC, Chili**


**Mohamed Zare Ernani**

**PhD fellow, Ghent University, Belgium**

**Deyanira Lobo**

**Universidad Central de Venezuela**

**Coordinator Andean Countries, CAZALAC, Chili**



## Why aridity indices????

To delineate zones prone to desertification  
*Those areas characterized by low rainfall and by high summer temperatures, so that the vegetation has little opportunity to restore after destruction by human impact or prolonged droughts.*

---- desertification????

---- aridity????

---- droughts????

# DESERTIFICATION

- The Earth Summit (UNCED, Rio de Janeiro, 1992) defined and  
The General Assembly of UN (UNCED, Paris, 1994) approved *the definition of DESERTIFICATION* :

*‘Desertification is land degradation in arid, semi-arid and dry sub-humid areas resulting from climatic variations and human activities’*



## DESERTIFICATION = drylands

- Drylands = arid , semi-arid, (dry)sub-humid
- 40 % of earth's surface
- 20% of earth's population
- Drylands are susceptible to 'human-induced soil degradation' and 'degradation of vegetation'



# Global extent of desertification

3 assessments

(mainly for political reasons)


Map 1/25.000.000 by FAO, UNESCO, WMO

## **1. UNCOD (1977):**

United Nations Conference on Desertification

Total area susceptible to desertification: 5281 Mha

Total area affected by desertification : 3970 Mha



# Global extent of desertification

- **2. GAP I (1984):**  
General Assessment of Progress  
(Plan of action to combat desertification)

questionnaire to 91 countries → failure  
UNEP used consultants

Distinction between: rainfed & irrigated cropland  
rangeland

 **STILL NO METHODOLOGY!!!!!!!!!!!!**



# Global extent of desertification

## ■ 3. GAP II (1992)

Used the Soil Data Base of GLASOD  
(Global Assessment of Soil Degradation)

Human induced soil degradation:

- |                               |         |
|-------------------------------|---------|
| - Degraded irrigated lands:   | 43 Mha  |
| - Degraded rainfed croplands: | 216 Mha |
| - Degraded rangelands:        | 757 Mha |

Rangelands with degraded vegetation: 2576 Mha

total 3592 Mha



# Global extent of desertification

## ■ Example of Africa

16% of surface:	water erosion
45% of surface:	wind erosion
25% of surface:	salinisation
61% of surface:	animal pressure
45% of surface:	population pressure



# ARIDITY VERSUS DROUGHT

## ■ ARIDITY

*Permanent* pluviometric deficit  
(long-term climatic phenomenon)

Linked to specific climatic data:

- strong insolation
- elevated temperatures
- low air humidity
- strong evapotranspiration

# ARIDITY VERSUS DROUGHT

## ■ DROUGHT

*Temporary* pluviometric deficit  
(short-term phenomenon)


Below average availability of natural water

**physical aspect:** below the long-term mean (normal)

**social aspect:** below expected volume to satisfy needs  
for agriculture, livestock, domestic use

DROUGHT is also: - annual/seasonal/monthly rainfall less than normal  
- reduced river flow

**DESERTIFICATION is commonly related to DROUGHT**



# DROUGHT

- Drylands are affected in an irregular manner by droughts

## Types of drought

- meteorological drought
- hydrological drought
- agricultural drought
- edaphic drought



# *Meteorological drought*

- Annual precipitation  $<$  average for one or several successive years

example: Sahel 1960s, 1970s, 1980s

**BUT!!!** *Average* is misleading because rain can be scattered or dry periods can alternate with periods of excessive rains.



## *Hydrological drought*

- Water resources used for industry, human, and animal consumption or support of agriculture (irrigation) is low.




# *Agricultural drought*

- Related to *rainfed agriculture*
- Related to ***soil moisture deficit*** during growing season

**Soil moisture deficit** can be determined by:

- precipitation
- PET
- soil moisture
- crop coefficient (moisture requirements)

Absolute annual or seasonal deficit of precipitation is  
not a good indicator →→better: rainfall distribution



## *Edaphic drought*

- →→ decrease of infiltrability of the soil
- →→ increase in runoff and erosion



# DELINEATING ARIDITY ZONES

- Based on **INDEX OF MOISTURE DEFICIT**

or

ARIDITY INDEX  $AI = P/PET$

# DETERMINATION OF PET

- 1) direct measurements using lysimeters

➡ **NO STANDARDIZATION**

- 2) empirical formulas

*Penman & Penman Monteith*

➡ **NEEDS LOT OF DATA:** solar radiation,  
wind velocity, relative humidity, temperature

- 3) relation between measured PET and two parameters

ex. *Thornthwaite*: mean monthly temperature and average number of  
daylight-hours/month

➡ **OVERESTIMATES** PET FOR **DRY** CONDITIONS


➡ **UNDERESTIMATES** PET FOR **MOIST** AND **COLD** CONDITIONS

# GLOBAL EXTENT OF ARIDITY ZONES

	<b>Aridity Index AI= P/PET</b>	<b>Million (ha)</b>	<b>% world land area</b>
<b>1.Cold</b>	<b>&gt; 0.65</b>	<b>1765.0</b>	<b>13.6</b>
<b>2.Humid</b>	<b>&gt; 0.65</b>	<b>5100.4</b>	<b>39.2</b>
<b>3.Dry sub-humid</b>	<b>0.50-0.65</b>	<b>1294.7</b>	<b>9.9</b>
<b>4.Semi-arid</b>	<b>0.20-0.50</b>	<b>2305.3</b>	<b>17.7</b>
<b>5.Arid</b>	<b>0.05-0.20</b>	<b>1569.1</b>	<b>12.1</b>
<b>6.Hyper-arid</b>	<b>&lt; 0.05</b>	<b>978.2</b>	<b>7.5</b>
<b>Drylands (3+4+5+6)</b>	<b>&lt; 0.65</b>	<b>6147.3</b>	<b>47.2</b>
<b>Susceptible Drylands (3+4+5)</b>	<b>0.05-0.65</b>	<b>5169.1</b>	<b>39.7</b>


# World drylands (Mha)

	Africa	Asia	Australia	Europe	North America	South America	World Total	%
<b>Hyper - arid</b>	<b><u>672</u></b>	<b>277</b>	<b>0</b>	<b>0</b>	<b>3</b>	<b>26</b>	<b>978</b>	<b>16</b>
<b>Arid</b>	<b>504</b>	<b><u>626</u></b>	<b>303</b>	<b>11</b>	<b>82</b>	<b>45</b>	<b>1571</b>	<b>26</b>
<b>Semi-arid</b>	<b>514</b>	<b><u>693</u></b>	<b>309</b>	<b>105</b>	<b>419</b>	<b>265</b>	<b>2305</b>	<b>37</b>
<b>Dry sub-humid</b>	<b>269</b>	<b><u>353</u></b>	<b>51</b>	<b>184</b>	<b>232</b>	<b>207</b>	<b>1296</b>	<b>21</b>
<b>Total</b>	<b><u>1959</u></b>	<b><u>1949</u></b>	<b>663</b>	<b>300</b>	<b>736</b>	<b>543</b>	<b>6150</b>	<b>100</b>
<b>% World Total</b>	<b><u>32</u></b>	<b><u>32</u></b>	<b>11</b>	<b>5</b>	<b>12</b>	<b>8</b>	<b>100</b>	
<b>% Total Global Land Area</b>	<b>13.1</b>	<b>13.0</b>	<b>4.4</b>	<b>2.0</b>	<b>4.9</b>	<b>3.6</b>	<b><u>41.0</u></b>	
<b>% Continent Area</b>	<b><u>66</u></b>	<b>46</b>	<b><u>75</u></b>	<b>32</b>	<b>34</b>	<b>31</b>	<b>41</b>	



# DRYLAND ZONES

- Based on climate and environmental attributes
  - **BUT!!!!** Dryland boundaries are neither static nor abrupt because of:
    - **1. high inter-annual variability in mean rainfall**
    - **2. occurrence of drought which may last for several years.**
- ⇒ Individual aridity zones do not represent homogeneous climates, either in the long term or during a particular time band
- ⇒ Dryland boundaries may not be defined in terms of natural vegetation or soil type because of human induced processes.



# **DRYLAND ZONES**

**Hyper-arid zones (environments)**

**Arid zones**

**Semi-arid zones**

**Dry sub-humid zones**




# Hyper-arid areas

Very limited rainfall

Highly variable rainfall: inter-annually (100%), monthly

Year-long periods without rainfall

⇒ true deserts ⇒ not prone to desertification  
⇒ very low biological productivity



# Arid zones


**< 200 mm in annual winter rainfall**

**< 300 mm in summer rainfall**

**Inter-annual variability 50-100%**

**Pastoralism is possible**

**Use of groundwater is highly  
susceptible to climate variability**



# Semi-arid zones

**Highly seasonal rainfall distribution**

**< 500 mm in winter rainfall regimes**


**< 800 mm in summer rainfall regimes**

**Inter-annual variability: 25-50%**

**Grazing of grassland**

**Sedentary agricultural activities are  
susceptible to seasonal and inter-annual  
deficiency**





# Dry sub-humid zones

High seasonal rainfall regimes

< 25% inter-annual variability

Rainfed agriculture

⇒ susceptible to degradation enhanced by seasonal rainfall, drought periods and increasing intensity of human use.

↪ **Dry sub-humid zones are included in the definition of desertification**



# ARIDITY INDICES

- Problem??
- - Data collection
- - The more parameters, the more errors




# **EVALUATION AND CLASSIFICATION OF CLIMATIC INDICES FOR YAZD REGION (IRAN)**

- **Mohammad Zare Ernani and Donald Gabriels**
- Department of Soil Management and Soil Care, International Centre for Eremology, Ghent University, Belgium





# PURPOSE:

- This study aims at comparing different climatic indices for evaluating the aridity and the rain aggressivity and rain distribution based on climatic data from 21 weather stations in the Yazd-Ardakan basin (Iran) and this for 5 to 48 successive years.



- Five aridity indices were used to assess the aridity in the basin:
  - De Martonne Aridity Index
  - Emberger Aridity Index
  - UNEP Aridity Index
  - Thornthwaite Classification
  - Gausse-Bagnouls Classification.
  
- For rain distribution and rain concentration
  - Modified Fournier Index (MFI)
  - Precipitation Concentration Index (PCI)





# 1. Aridity index of De Martonne

$$I_M = \frac{P}{t + 10}$$

$P$  = Annual average rainfall in mm.

$t$  = Annual average temperature in degrees centigrade.

Climate Type	Aridity Index
Arid	0-10
Semi-arid	10-20
Mediterranean	20-24
Semi-humid	24-28
Humid	28-35
Very Humid	35-55
Extremely Humid	>55

# 2. Aridity index of Emberger

$$I_E = \frac{100 \times P}{M^2 - m^2}$$

$M$  = Average temperature of the hottest month in degrees centigrade.

$m$  = Average temperature of the coldest month in degrees centigrade.

$P$  = Annual average rainfall in mm.

### 3. UNEP aridity index (P/ETP)

- $ET_o = 16 \times Nm \left( \frac{10 \times T_m}{I} \right)^a$
- $T_m$  = mean monthly temperature
- $N_m$  = adjustment factor related to hours of daylight
- Heat Index or  $I = \sum (T_m / 5)^{1.514}$  for  $m = 1 \dots 12$
- $a = 6.75 \times 10^{-7} \times I^3 - 7.71 \times 10^{-5} \times I^2 + 1.792 \times 10^{-2} \times I + 0.49239$

$$I = \sum_1^{12} (T_m / 5)^{1.514}$$


Index	Class
$P/ETP < 0.03$	hyper-arid zone
$0.03 < P/PET < 0.2$	arid zone
$0.2 < P/PET < 0.5$	semi-arid zone
$P/PET > 0.5$	humid zone

### 4. Thornthwaite classification

$$PE_{index} = \sum_1^{n=12} 115 \times \left( \frac{P}{T - 10} \right)^{10/9}$$

- $P$  = monthly precipitation in inches;
- $T$  = temperature in °F; and  $n$  = months = 12.


PE Index	Climate
More than 128	Wet
64127	Humid
3263	Sub-Humid
1631	Semiarid
Less than 16	Arid




## 5. Gaussen-Bagnouls classification method


- combination of average monthly temperature and total rainfall
- gives more precise climatic classification
- easily climatic identification by determining separately the numbers of dry and wet months


$P > 3T$  → Humid  
 $3T > P > 2T$  → Semi-humid  
 $P < 2T$  → Arid






## Yazd Precipitation





# YAZD RAINFALL MOVING AVERAGE



◆ Series 1

— 7 per. Mov. Avg. (Series 1)

— 61.58



## 6. Precipitation concentration index

- to estimate the temporal variability of monthly rainfall


$$PCI = 100 \frac{\sum p_i^2}{P^2}$$

$p$  = monthly precipitation


$P$  = annual precipitation

PCI	Concept
8.3 – 10	uniform
10 – 15	Moderately seasonal
15 – 20	seasonal
20 – 50	Highly seasonal
50 – 100	Irregular

<b>STATION</b>	<b>P/ETP</b> (UNEP method)	<b>PE Index</b> (Thornthwaite classification)	<b>I<sub>EMBERGER</sub></b>	<b>I<sub>De Martonne</sub></b>
<b>ANAR</b>	<b>0.08</b>	<b>4.72</b>	<b>5.85</b>	<b>2.91</b>
<b>ARDAKAN</b>	<b>0.06</b>	<b>3.81</b>	<b>3.90</b>	<b>2.22</b>
<b>ASHKZAR</b>	<b>0.06</b>	<b>3.79</b>	<b>4.00</b>	<b>2.30</b>
<b>BAFGH</b>	<b>0.04</b>	<b>3.16</b>	<b>3.35</b>	<b>1.74</b>
<b>ESFANDABAD</b>	<b>0.07</b>	<b>3.40</b>	<b>4.39</b>	<b>2.41</b>
<b>KAVIRSLAHKOOH</b>	<b>0.06</b>	<b>4.81</b>	<b>4.38</b>	<b>2.55</b>
<b>KHORBIABANAK</b>	<b>0.06</b>	<b>4.07</b>	<b>4.64</b>	<b>2.52</b>
<b>NAEEN</b>	<b>0.10</b>	<b>6.44</b>	<b>7.28</b>	<b>3.65</b>
<b>NASRABAD</b>	<b>0.19</b>	<b>10.30</b>	<b>14.53</b>	<b>6.05</b>
<b>ROBATEPOSHT</b>	<b>0.10</b>	<b>5.79</b>	<b>8.27</b>	<b>3.76</b>
<b>SAGHAND</b>	<b>0.06</b>	<b>5.76</b>	<b>4.05</b>	<b>2.12</b>
<b>VARZANEH</b>	<b>0.08</b>	<b>4.72</b>	<b>5.24</b>	<b>2.89</b>
<b>YAZD</b>	<b>0.05</b>	<b>2.92</b>	<b>3.99</b>	<b>2.11</b>



Station	PCI <sub>1</sub>	PCI <sub>2</sub>	P <sub>mean</sub>	PERIOD
ABARKUH	29.87	14.3	60.1	1967-1995
ARDEKAN	28.36	16.54	55.56	1966-1990
ASHKZAR	24.98	16.07	67.65	1978-1995
BAJGAN	26.66	15.05	235.61	1966-1995
DEHSHIR	26.67	14.34	95.87	1967-1995
GHOTROOM	23.87	15.62	140.8	1966-1995
HAJIABAD	26.66	14.2	77.36	1966-1995
KHARANAGH	22.05	14.98	129.82	1966-1995
KHOOR BIABANAK	28.43	16.61	88.02	1986-1995
MAZRAEH NOW	28.21	16.21	98.54	1967-1995
NASRABAD	21.01	15.22	211.97	1967-1995
ROBATPOSHTE	25.15	15.55	131.75	1967-1989
TAFT	23.57	15.75	131.68	1966-1994*
VARZANEH	28.21	13.76	73.01	1958-1995
ANAR	29.64	16.16	84.90	1986-2000
BAFGH	30.15	18.44	58.10	1993-2000
HOJATABAD	21.33	15.74	155.48	1967-1985
KAVIR SIYAHKOOH	31.59	19.62	73.70	1988-2000
NAEEN	25.88	14.74	99.95	1969-2000
SAGHAND	28.23	15.74	67.94	1967-2000
YAZD	29.80	15.29	61.58	1953-2000



## Dry and humid periods

$$LP_D = \text{N}^\circ \text{ of dry months} = P < 0,5 ET_0$$

$$LP_R = \text{N}^\circ \text{ of rainy months} = P > 0,5 ET_0$$

where: :  $LP_D$ : length of the water shortage period  
 $LP_R$ : length of the water surplus period

## Water shortage

$$DH = \sum_1^{12} (P - ET_0)$$

DH = yearly water shortage (mm)  
P = monthly precipitation (mm)



# MAP of Aridity Zones in Latin America

**Project CAZALAC**  
(**Centro del Agua para Zonas  
Áridas y Semiáridas de  
América LATina y El Caribe**),  
**La Serena, Chili**


**Deyanira Lobo Luján**  
**Universidad Central de Venezuela**

**Coordinator Andean Countries**

# Penman-Monteith-FAO

$T_m$ , monthly mean temperature ( $^{\circ}\text{C}$ )  
 $R_s$ , solar radiation in  $\text{calcm}^{-2}\text{d}^{-1}$   
 $HR$ , monthly mean relative humidity (%)  
 $U_2$ , wind speed in  $\text{ms}^{-1}$

Characteristics of a hypothetical reference crop  
(green grass of 0.12 m high with an albedo of 0.23)



$r_a$ : aerodynamic resistance

$r_s$ : surface resistance of the green grass =  $70 \text{ s/m}$

## Penman-Montheith-FAO

$$ET_0 = \frac{0.408\Delta(R_n - G) + \gamma \frac{900}{T + 273} u_2 (e_s - e_a)}{\Delta + \gamma(1 + 0.34u_2)}$$

Where:

**ET<sub>0</sub>** baseline evapotranspiration [mm day<sup>-1</sup>],

**R<sub>n</sub>** net radiation on the crop surface [MJ m<sup>-2</sup> day<sup>-1</sup>],

**G** heat flow density in the soil [MJ m<sup>-2</sup> day<sup>-1</sup>],

**T** average daily temperature at a 2 m height [°C],

**u<sub>2</sub>** wind speed at a 2 m height [m s<sup>-1</sup>],

**e<sub>s</sub>** saturation vapor pressure [kPa],

**e<sub>a</sub>** current vapor pressure [kPa],

**e<sub>s</sub> - e<sub>a</sub>** saturation vapor pressure gap [kPa],

**Δ** slope of the saturation vapor pressure line in function of temperature [kPa °C<sup>-1</sup>],

**γ** psychrometric coefficient [kPa °C<sup>-1</sup>].



## Protocol to calculate the baseline Evapotranspiration by using the FAO/Penman-Monteith equation

**Step 1. Calculation of Net Radiation, Rn (MJ/m<sup>2</sup> day) from global solar radiation, RG (Cal/cm<sup>2</sup> day)**

$$R_n = RG * .0419 * .8$$

Factor 0.0419 converts cal/cm<sup>2</sup> day into MJ/m<sup>2</sup> day

Factor 0.8 is the Rn/RG quotient for a vegetated area with a good water supply

**Step 2. Calculation of air vapor pressure at saturation, es (kPa)**

$$e_s = 0.707 * \text{EXP}(.05979 * T_a)$$

T<sub>a</sub> is the average air temperature (°C)

**Step 3. Calculation of air saturation shortage Ds(kPa)**

$$D_s = e_s * (1 - HR / 100)$$

HR is relative humidity in %



**Step 4. Calculation of the saturation vapor pressure curve slope,  $\Delta$  (kPa/°C)**

$$\begin{aligned}TETA &= (T_a + 237.3)^2 \\ALFA &= 17.27 * T_a / (T_a + 237.3) \\ \Delta &= 4098 * (.6108 * \text{EXP}(ALFA)) / TETA\end{aligned}$$

**Step 5. Calculation of advective contribution**

$$\text{Adv} = (\gamma * 900 * U * D_s) / (T_a + 273)$$

$$\gamma = .066 \text{ (kPa/°C)}$$

U = wind speed in m/s


D<sub>s</sub> = saturation shortage (kPa)

**Step 6. Calculation of the radioactive contribution**

$$\text{Rad} = 0.408 * \Delta * R_n$$

**Step 7. Calculation of denominator (resistance to vapor diffusion in the limit layer)**

$$D_n = \Delta + .066 * (1 + .34 * U)$$



**Step 8. Calculation of the radioactive component of ETo**

$$ETRAD = Rad / Dn$$

**Step 9. Calculation of the advective component of ETo**

$$ETADV = Adv / Dn$$

**Step 10. Calculation of ETo**

$$ETo = ETRAD + ETADV$$

## PRELIMINARY MAP OF ARID ZONES OF SOUTH AMERICA DEMO VERSION





***Gezondheid !  
Cheers!***

***Saude!***

***Salud!***

***Santé!***



**Thank you**

***Merçi***

**Gracias**

**Obrigado**

***Dank u***

